

An uncountable union of line segments with null two-dimensional measure

Parker Kuklinski

October 2023

Contents

Image copy	3
Text copy with conversion errors	12
Abstract	12
1. Introduction	12
2. No lower bound	13
3. Parallelogram formulation	15
4. Proving the conjectures	16
References	19

Image copy

An uncountable union of line segments with null two-dimensional measure

Parker Kuklinski

October 2023

Abstract

In this paper we construct an uncountable union of line segments T which has full intersection with the sets $(\{0\} \times [0, 1]) \cup (\{1\} \times [0, 1]) \subset \mathbb{R}^2$ but has null two-dimensional measure. Further results are proved on the decay rate of $\mu(T)$ if the line segments comprising T are replaced with increasingly fine approximations by parallelograms.

1 Introduction

In [1], Kaczynski proves a variety of results placing restrictions on possible boundary functions of a half-plane function with some property. The last result is different; he constructs a measurable half plane function f with a nonmeasurable boundary function. Crucial to this construction is a measure theoretic generalization of the trapezoid formula which we document here. For the remainder of the paper, let $X_0 = \{(x, y) : y = 0\}$ and $X_1 = \{(x, y) : y = 1\}$, let μ refer to Lebesgue measure in the contextually obvious dimension (we will distinguish μ_1 and μ_2 to refer to one and two-dimensional Lebesgue measure respectively when necessary), and let μ^* refer to outer measure.

Definition 1 A trapezoid $T = \cup \ell$ is a disjoint union of line segments with each segment having its two endpoints on X_0 and X_1 (i.e. $\ell = \{(1-t)x_0 + tx_1, t\} : t \in [0, 1]\}$).

If T is a trapezoid, let $T_0 = T \cap X_0$ and $T_1 = T \cap X_1$. In [1], Kaczynski proved the following result:

Theorem 1 If T is a trapezoid, then $\mu_2^*(T) = (\mu_1^*(T_0) + \mu_1^*(T_1))/2$.

This theorem was used to construct a collection of non-intersecting paths (not lines) γ_x corresponding to each point $x \in [0, 1]$ such that for each path $\lim_{t \rightarrow 0} \gamma_x(t) = (x, 0)$ but the union satisfies $\mu(\cup_{x \in [0, 1]} \gamma_x) = 0$. This construction immediately leads to the creation of a measurable half plane function with non-measurable boundary function [1]. We depict the situation in Figure 1.

One may wonder the extent to which this result holds if we omit the restriction that the lines of a trapezoid are not disjoint. To this end, we amend the trapezoid definition:

Definition 2 An unrestricted trapezoid $T = \cup \ell$ is a union (not necessarily disjoint) of line segments with each segment having its two endpoints on X_0 and X_1 (i.e. $\ell = \{(1-t)x_0 + tx_1, t\} : t \in [0, 1]\}$).

In [1], Kaczynski proves that there is no sensible upper bound in terms of $\mu_1(T_0)$ and $\mu_1(T_1)$ on the two dimensional measure of an unrestricted trapezoid.

Theorem 2 There exists an unrestricted trapezoid T with $\mu_1(T_0) = \mu_1(T_1) = 0$ and $\mu_2(T) = \infty$.

Proof: Let M be a residual set of measure zero (e.x. let x_k be an enumeration of rationals and $M = \bigcap_{n=1}^{\infty} \bigcup_{k=1}^{\infty} (x_k - \frac{1}{2^{k+n}}, x_k + \frac{1}{2^{k+n}})$). For any point (x, y) ($y \in (0, 1)$) and line ℓ passing through (x, y) with angle θ with the x -axis, let $F_0(\theta) = (x - y \cot(\theta), 0)$ and $F_1(\theta) = (x + (1 - y) \cot(\theta), 1)$ denote the intersections $\ell \cap X_0$ and $\ell \cap X_1$ respectively. F_0 and F_1 are homeomorphisms from $(0, \pi)$ onto X_0 and X_1 respectively, so $F_0^{-1}(M)$ and $F_1^{-1}(M)$ are both residual sets of measure zero. Since the intersection of residual




Figure 1: Plot of $\cup \gamma_x$ for $x \in [0, 1]$ where $\gamma_{x_1} \cap \gamma_{x_2} = \emptyset$, $\lim_{t \rightarrow 0} \gamma_x(t) = (x, 0)$, but $\mu(\cup_{x \in [0, 1]} \gamma_x) = 0$. Given a nonmeasurable function $\varphi(x)$, if we define $f : (\mathbb{R} \times (0, \infty)) \rightarrow \mathbb{R}$ as $f(\gamma_x) = \varphi(x)$ and vanishing elsewhere, then f is measurable in the upper half plane but has a nonmeasurable boundary function φ .

sets is nonempty, for every (x, y) with $y \in (0, 1)$ we can choose an angle $\alpha = F_0^{-1}(M) \cap F_1^{-1}(M)$ such that the line ℓ intersecting (x, y) with angle α intersects $X_0 \cap M$ and $X_1 \cap M$. Let the unrestricted trapezoid T be the union of all such lines so that $\mu_1(T_0) = \mu_1(T_1) = \mu_1(M) = 0$ and $\mu_2(T) = \mu_2(\{(x, y), y \in (0, 1)\}) = \infty$. \square .

2 No lower bound

In [1], Kaczynski poses the opposite question as an open problem, asking if there is a lower bound on the two-dimensional measure of an unrestricted trapezoid in terms of $\mu(T_0)$ and $\mu(T_1)$. We will prove the following theorem indicating no lower bound exists:

Theorem 3 *There exists an unrestricted trapezoid T with $\mu_1(T_0) = \mu_1(T_1) = \infty$ and $\mu_2(T) = 0$.*

Since it suffices to find an unrestricted trapezoid with measure zero and sets $\mu(T_0) = \mu(T_1) = \mu([0, 1]) = 1$, we can formulate the question a different way:

Proposition 1 *There exists a bijection $f : [0, 1] \rightarrow [0, 1]$ such that, for the set*

$$S = \bigcup_{t \in [0, 1]} \bigcup_{x \in [0, 1]} ((1-t)x + tf(x), t),$$

we have $\mu(S) = 0$.

It is clear that Theorem 3 is a corollary of Proposition 1.

It is perhaps not immediate that such a function should exist. Any monotone f induces a standard trapezoid which abides by Theorem 1. The choice $f(x) = 1 - x$ gives measure $\mu(S) = 1/2$, and other decreasing functions cannot be massaged to induce a smaller measure. We will instead draw inspiration




Figure 2: Examples of the fractal construction of an unrestricted trapezoid with null measure. (Left) The base pattern induced by f_1 (Right) Illustration of unrestricted trapezoid induced by f .

from fractal geometry [2]; the particular iterative scheme we choose will create one-dimensional slices in x similar to the Cantor middle third set. In Figure 2, the basic pattern is represented by the piecewise function

$$f_1(x) = \begin{cases} x & x \in [0, \frac{1}{3}) \\ x + \frac{1}{3} & x \in [\frac{1}{3}, \frac{2}{3}) \\ x - \frac{1}{3} & x \in [\frac{2}{3}, 1] \end{cases}$$

otherwise we divide $[0, 1]$ into three subintervals and connect them via parallelograms where f_1 essentially swaps the second and the third. In this case, $\mu(S) = 5/6$. In each subsequent iteration we embed an affine transformed version of the base pattern in each parallelogram, thus reducing the total measure of the unrestricted trapezoid (we will prove this limits to zero as in Proposition 1).

Since this procedure iteratively divides subintervals of $[0, 1]$ in 3, it is perhaps not surprising that this sequence of functions $\{f_k\}$ has a natural representation on base 3 fractions, namely

$$f_k(0.x_1x_2x_3, \dots) = 0.y_1y_2\dots y_kx_{k+1}x_{k+2}\dots$$

where $y_j = 2x_j \pmod{3}$. Let $f = \lim_{k \rightarrow \infty} f_k$ such that for input $x \in [0, 1]$, f replaces every 1 in its base 3 expansion with a 2 and vice-versa. We will prove that this function satisfies proposition 1.

Before we prove proposition 1, we will state a result from [3] on modified Cantor sets:

Proposition 2 Let C_t be a modified Cantor set with representation

$$C_t = \left\{ \sum_{k=1}^{\infty} \frac{x_k}{3^k} : x \in \{0, 1, t\} \right\}.$$

Then the following holds:

$$\mu_1(C_t) = \begin{cases} 1/q & t = p/q \text{ in lowest terms, } p + q \equiv 0 \pmod{3} \\ 0 & \text{otherwise} \end{cases}$$

We are now ready to prove Proposition 1.

Proof of Proposition 1: Let us define f acting on base 3 fractions as

$$f_k(0.x_1x_2x_3, \dots) = 0.y_1y_2y_3\dots$$

where $y_j = 2x_j \pmod{3}$. Then the linear combination $(1-t)x + tf(x)$ satisfies

$$S_t = \bigcup_{x \in [0,1]} ((1-t)x + tf(x)) = \left\{ \sum_{k=1}^{\infty} \frac{x_k}{3^k} : x \in \{0, 1+t, 2-t\} \right\}.$$

Since S_t is just a scaled copy of $C_{(2-t)/(1+t)}$, by Proposition 2 we have the measure

$$\mu_1(S_t) = \begin{cases} (1+t)/q & (2-t)/(1+t) = p/q \text{ in lowest terms, } p+q \equiv 0 \pmod{3} \\ 0 & \text{otherwise} \end{cases}$$

Since $\mu_1(S_t)$ takes on nonzero value at only countably many t , the result follows from Fubini's theorem. \square

3 Parallelogram formulation

Over 30 years after [1], Kaczynski would outline a reformulation of the problem, offering three conjectures on the matter [4]. Rather than conceiving the unrestricted trapezoid as a union of lines, Kaczynski considered unions of increasingly thinner parallelograms.

Definition 3 Let $P_{j,k}^n = \{(x, y) \in \mathbb{R}^2 : nx + (k-j)y \geq k-1, nx + (k-j)y \leq k, y \in [0, 1]\}$ be a parallelogram with parallel sides $[(j-1)/n, j/n] \times \{0\}$ and $[(k-1)/n, k/n] \times \{1\}$. For $\sigma \in \text{Sym}(n)$, we say that the unrestricted trapezoid T_σ^n is given by the union $T_\sigma^n = \bigcup_{j=1}^n P_{j, \sigma(j)}^n$.

In this way, the unrestricted trapezoid associated with f_1 above can be expressed as $T_{(2\ 3)}^3$.

The following conjectures center around properties of the smallest unrestricted trapezoid of n parallelograms of width $1/n$, namely $\alpha(n) = \min_{\sigma \in \text{Sym}(n)} \mu_2(T_\sigma^n)$.

Conjecture 1 $\lim_{n \rightarrow \infty} \alpha(n) = 0$.

Kaczynski claimed to have proved this. We will show that this follows from Proposition 1 in the following section.

Conjecture 2 There exists constants $c_\pm > 0$ such that $\frac{c_-}{\log n} < \alpha(n) < \frac{c_+}{\log n}$.

Kaczynski claimed to be able to prove that c_- exists and also claimed a weaker upper bound in the form of $\alpha(n) < c_+ (\log(\log n))^2 / \log n$. We will prove a different upper bound.

Conjecture 3 The sequence $\alpha(n)$ decreases monotonically.

This was left as an open problem in [4].

4 Proving the conjectures

To prove conjectures from the previous section, we first connect projections of the Sierpinski gasket [3] to our modified Cantor sets.

Definition 4 We define the Sierpinski gasket $\mathcal{G} \subset \mathbb{R}^2$ as

$$\mathcal{G} = \left\{ \sum_{k=1}^{\infty} \frac{x_k}{3^k} : x \in \{(0, 0), (2, 0), (0, 2)\} \right\}.$$

We define the n^{th} partial Sierpinski gasket $\mathcal{G}_n \supset \mathcal{G}$ similarly. Let $G_n = \{\sum_{k=1}^n \frac{x_k}{3^k} : x \in \{(0, 0), (2, 0), (0, 2)\}\}$ and let $\Delta(a, b, c)$ be the area bounded by the triangle with vertices $a, b, c \in \mathbb{R}^2$.

$$\mathcal{G}_n = \bigcup_{x \in G_n} \Delta\left(x, x + \left(\frac{1}{3^n}, 0\right), x + \left(0, \frac{1}{3^n}\right)\right)$$

Let the projection operator $\text{proj}_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}$ map points $(x, y) \in \mathbb{R}^2$ onto a line through the origin with angle θ to the x -axis such that $\text{proj}_\theta : (x, y) \mapsto x \cos \theta + y \sin \theta$. We are interested in the sets $\text{proj}_\theta(\mathcal{G}_n)$; more generally there is an active field of research surrounding linear projections of fractals in \mathbb{R}^2 [5]. One of the major subjects of interest is the following:

Definition 5 The Favard distance of a compact set $E \subset \mathbb{R}^2$ is given by

$$\text{Fav}(E) = \frac{1}{\pi} \int_0^\pi \mu(\text{proj}_\theta(E)) d\theta$$

While it is perhaps obvious that $\text{Fav}(\mathcal{G}) = 0$ from proposition 2, a result by Bond and Volberg [6] gives a decay bound on the Favard distances of partial Sierpinski gaskets.

Proposition 3 There exist constants $C, p > 0$ such that $\text{Fav}(\mathcal{G}_n) \leq Cn^{-p}$.

A later result by Bond and Volberg proves that $p > 1/14$ [7].

The rest of this section centers around utilizing proposition 3 to prove the conjectures from section 3. To do this, we must compare $\text{proj}_\theta(\mathcal{G}_n)$ to the modified Cantor sets from Proposition 2 and the horizontal slices of the unrestricted trapezoid. Let us define partial variants of these sets:

Definition 6 Let $D_n(a, b, c) = \{\sum_{k=1}^n \frac{x_k}{3^k} : x_k \in \{a, b, c\}\}$. If $T_\sigma^{3^n}$ is the unrestricted trapezoid associated with the n^{th} iteration induced by swapping $1 \leftrightarrow 2$ in a base 3 expansion (i.e. if $x - 1 = x_n x_{n-1} \dots x_1$ is an n -digit base 3 number, then $\sigma(x - 1) + 1 = y_n y_{n-1} \dots y_1$ where $y_k = 2x_k \pmod{3}$), then we write $S_t^{(n)} = (\mathbb{R} \times \{t\}) \cap T_\sigma^{3^n}$, the slice of $T_\sigma^{3^n}$ at $y = t$, as

$$S_t^{(n)} = \bigcup_{x \in D_n(0, 1+t, 2-t)} \left[x, x + \frac{1}{3^n} \right]$$

We similarly define the n^{th} partial modified Cantor set as

$$C_t^{(n)} = \bigcup_{x \in D_n(0, 1, t)} \left[x, x + \frac{1}{3^n} \right]$$

This allows us to make the following comparison:

Lemma 1 For $t \in [0, 1]$ and $\varphi(t) = \tan^{-1} \left(\frac{2-t}{1+t} \right)$, we have

$$\mu(S_t^{(n)}) \leq (1+t) \mu(\text{proj}_{\varphi(t)}(\mathcal{G}_n))$$

Proof: We can show that $\mu(S_t^{(n)}) \leq (1+t) \mu(C_{(2-t)/(1+t)}^{(n)})$ for $t \geq 0$ since

$$\begin{aligned} S_t^{(n)} &= \bigcup_{x \in D_n(0, 1+t, 2-t)} \left[x, x + \frac{1}{3^n} \right] \subset \bigcup_{x \in D_n(0, 1+t, 2-t)} \left[x, x + \frac{1+t}{3^n} \right] \\ &= \bigcup_{x \in D_n(0, 1, \frac{2-t}{1+t})} (1+t) \left[x, x + \frac{1}{3^n} \right] = (1+t) C_{(2-t)/(1+t)}^{(n)} \end{aligned}$$

The result then follows from noticing that for $\theta \leq \frac{\pi}{4}$, we have $\text{proj}_\theta(\mathcal{G}_n) = C_{\tan \theta}^{(n)}$. \square

Combining these results leads us to a similar polynomial bound on the decay of the unrestricted trapezoids induced by the iterative scheme described in Definition 6.

Proposition 4 Let σ be the permutation on base 3 numbers as defined in Definition 6. Then there exist constants $C, p > 0$ such that $\mu(T_\sigma^{3^n}) \leq Cn^{-p}$.

Proof: By Fubini's theorem, The measure of $T_\sigma^{3^n}$ is given by integrating over the measures of the horizontal slices

$$\mu_2(T_\sigma^{3^n}) = \int_0^1 \mu(S_t^{(n)}) dt$$

From Lemma 1, this integral can be bound by an integral over projections of the Sierpinski gasket:

$$\int_0^1 \mu(S_t^{(n)}) dt \leq \int_0^1 (1+t) \mu(\text{proj}_{\varphi(t)}(\mathcal{G}_n)) dt$$

By the change of variables $\theta = \varphi(t)$, we can write:

$$\int_0^1 (1+t) \mu(\text{proj}_{\varphi(t)}(\mathcal{G}_n)) dt = \int_{\tan^{-1}(1/2)}^{\tan^{-1}(2)} \frac{9(\tan^2 \theta + 1)}{(\tan \theta + 1)^3} \mu(\text{proj}_\theta(\mathcal{G}_n)) d\theta$$

Placing an estimate on the trigonometric portion of the integral we can write

$$\int_{\tan^{-1}(1/2)}^{\tan^{-1}(2)} \frac{9(\tan^2 \theta + 1)}{(\tan \theta + 1)^3} \mu(\text{proj}_\theta(\mathcal{G}_n)) d\theta \leq \frac{10}{3} \int_{\tan^{-1}(1/2)}^{\tan^{-1}(2)} \mu(\text{proj}_\theta(\mathcal{G}_n)) d\theta$$

Since Lebesgue measure is nonnegative, we can extend the limits of integration to bound the measure of $T_\sigma^{3^n}$ by a factor of the associated Favard distance

$$\mu(T_\sigma^{3^n}) \leq \frac{10\pi}{3} \text{Fav}(\mathcal{G}_n)$$

Proposition 3 completes the proof. \square

This result translates to a logarithmic bound on $\alpha(n)$ *only* for $n = 3^m$ where $m \in \mathbb{N}$. If n has the base 3 representation $n = x_k x_{k-1} \dots x_0$, we can define σ_n as in Figure 3 such that the first $x_k 3^k$ intervals map to x_k copies of $T_\sigma^{3^k}$, the next $x_{k-1} 3^{k-1}$ intervals map to x_{k-1} copies of $T_\sigma^{3^{k-1}}$, etc. Using this scheme, we can prove a weaker upper bound on Conjecture 2 (consequently proving Conjecture 1 as well):

Theorem 4 *There exist constants $C, p > 0$ such that $\alpha(n) < C/(\log n)^p$.*

Before proving this theorem, we supply a technical lemma:

Lemma 2 *For $p \in (0, 1)$, there exists a constant C such that*

$$\int_0^n e^x x^{-p} dx \leq C e^n n^{-p}$$

Proof: Let us make the change of variables $t = 1 - x/n$ such that this integral can be transformed into

$$\int_0^n e^x x^{-p} dx = \frac{e^n}{n^{p-1}} \int_0^1 \frac{e^{-nt}}{(1-t)^p} dt$$

For any constant $C > 1$, we can find t_0 dependent only on C, p such that $pt + 1 > (1-t)^{-p}$ for $t \in (0, t_0)$. This allows us to split the integral

$$\int_0^1 \frac{e^{-nt}}{(1-t)^p} dt \leq \int_0^{t_0} C(pt+1)e^{-nt} dt + \int_{t_0}^1 \frac{e^{-nt_0}}{(1-t)^p} dt$$

Both of these integrals have closed form expressions, the leading asymptotic term $1/n$ arising from the left integral, thus completing the proof. \square

Proof of Theorem 4: It is clear that $\alpha(n) \leq \mu(T_{\sigma_n}^n)$. Let $n = x_k x_{k-1} \dots x_0$ be a base 3 integer representation. Then the following holds:

$$\mu(T_{\sigma_n}^n) = \frac{1}{n} \sum_{j=0}^k x_j 3^j \mu(T_\sigma^{3^j})$$




Figure 3: Examples of parallelgram-defined unrestricted trapezoids T_σ^n where σ is as outlined in the proof of Theorem 4 (*Left*) $n = 1000$ which has base-3 representation 1101001 (*Right*) $n = 2000$ which has base-3 representation 2202002.

Each digit must satisfy $x_j \leq 2$, and by Proposition 4 there exist constants $C, p > 0$ such that $\mu(T_\sigma^{3^j}) \leq Cj^{-p}$, so we may write

$$\frac{1}{n} \sum_{j=0}^k x_j 3^j \mu(T_\sigma^{3^j}) < \frac{C}{n} \sum_{j=1}^{\lfloor \log_3 n \rfloor} 3^j j^{-p}$$

Since the function $f(x) = 3^x x^{-p}$ is increasing for $x > p/\log_3(x)$, we can bound this sum by the integral

$$\frac{C}{n} \sum_{j=1}^{\lfloor \log_3 n \rfloor} 3^j j^{-p} \leq \frac{C}{n} \int_0^{\log_3 n+1} 3^x x^{-p} dx$$

By making the change of variables $x = u \log_3 e$, we can write

$$\frac{C}{n} \int_0^{\log_3 n+1} 3^x x^{-p} dx = \frac{C(\log_3 e)^{1-p}}{n} \int_0^{\log 3n} e^u u^{-p} du$$

The result follows from applying Lemma 2 and collecting constants. \square

5 Application to Cluster Sets

While Theorem 1 assisted in constructing a measurable half-plane function with nonmeasurable boundary function in [1], the open questions from [1] and [4] answered by Propositions 1 and 4 were also initially conceived to address problems related to boundary functions and cluster sets. We state the relevant material here for completeness.

Definition 7 Given a function defined on the open unit disk $f : \mathbb{D} \rightarrow \mathbb{C}$ and a Jordan arc γ with an endpoint on the boundary of the disk, we say that $C(f, \gamma)$ is the cluster set of f on γ where

$$C(f, \gamma) = \{w \in \mathbb{C} : \exists \{z_n\} \text{ s.t. } z_n \in \gamma, |\lim z_n| = 1, \lim f(z_n) = w\}$$

Definition 8 Let $\gamma_1, \gamma_2, \gamma_3 \subset \mathbb{D}$ be Jordan arcs with a similar endpoint z on the unit circle. If, for a disk function f , the intersection satisfies $C(f, \gamma_1) \cap C(f, \gamma_2) \cap C(f, \gamma_3) = \emptyset$, then f has the three-arc property at z . If each $\gamma_1, \gamma_2, \gamma_3$ is a union of rectilinear segments, then we say f has the three-segment property at z .

In [1], a version of conjecture 1 was suggested to assist in answering the following open question from [8] (presented as conjecture):

Conjecture 4 *There exists a continuous function in \mathbb{D} having the three-segment property at each point of a set of positive measure or second category on $|z| = 1$.*

Weaker versions of this conjecture have been proved. Jarník [9] gave an example of a function having the three-segment property at uncountably many points. Piranian was cited in [8] as having proved the existence of a continuous disk function with the three-arc (not segment) property. Bagemihl later proved the existence of a normal meromorphic (not continuous) disk function with the three-segment property [10].

References

- [1] Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.
- [2] Kenneth J. Falconer. *Techniques in fractal geometry*, volume 3. Chichester: Wiley, 1997.
- [3] Richard Kenyon. Projecting the one-dimensional sierpinski gasket. *Israel Journal of Mathematics*, 97:221–238, 1997.
- [4] Yahoo! News. The unabomber letters: A yahoo news special report, 2016. <https://www.yahoo.com/entertainment/the-unabomber-letters---a-yahoo-news-special-report-170846210.html>.
- [5] Izabella Laba. Recent progress on favard length estimates for planar cantor sets. In *Operator-Related Function Theory and Time-Frequency Analysis*, pages 117–145. Cham: Springer International Publishing.
- [6] Matthew Bond and Alexander Volberg. The power law for buffon’s needle landing near the sierpinski gasket. arXiv preprint arXiv:0911.0233, 2009.
- [7] Matthew Bond and Alexander Volberg. Buffon needle lands in ϵ -neighborhood of a 1-dimensional sierpinski gasket with probability at most $|\log \epsilon|^{-c}$. *Comptes Rendus. Mathématique*, 348(11-12):653–656, 2010.
- [8] Frederick Bagemihl, George Piranian, and G. S. Young. Intersections of cluster sets. *Bulletin of the Polytechnic Institute of Iași*, 5(9):29–34, 1959.
- [9] Vojtěch Jarník. Sur les fonctions de deux variables réelles. *Fundamenta Mathematicae*, 27:147–150, 1936.
- [10] Frederick Bagemihl. The three-arc and three-separated-arc properties of meromorphic functions. *Nagoya Mathematical Journal*, 53:137–140, 1974.

Text copy with conversion errors

The version below contains many errors where symbols haven't been converted to html correctly, but it may be useful for word searching, such as being able to click the hyperlinked references at the end of the document and helping this document show up in searches.

Abstract

In this paper we construct an uncountable union of line segments T which has full intersection with the sets $(\{0\} \times [0,1]) \times \mathbb{R}^2$ and $(\{1\} \times [0,1]) \times \mathbb{R}^2$ but has null two-dimensional measure. Further results are proved on the decay rate of $\mu(T)$ if the line segments comprising T are replaced with increasingly fine approximations by parallelograms.

1. Introduction

In¹, Kaczynski proves a variety of results placing restrictions on possible boundary functions of a half-plane function with some property. The last result is different; he constructs a measurable half plane function f with a nonmeasurable boundary function. Crucial to this construction is a measure theoretic generalization of the trapezoid formula which we document here. For the remainder of the paper, let $X_0 = \{(x,y) : y = 0\}$ and $X_1 = \{(x,y) : y = 1\}$, let μ refer to Lebesgue measure in the contextually obvious dimension (we will distinguish μ_1 and μ_2 to refer to one and two-dimensional Lebesgue measure respectively when necessary), and let μ^* refer to outer measure.

Definition 1 A trapezoid $T = \bigotimes_{t \in [0,1]} \{(x,y) : (1-t)x_0 + tx_1, y = t\}$ is a disjoint union of line segments with each segment having its two endpoints on X_0 and X_1 (i.e. $\bigotimes = \{((1-t)x_0 + tx_1, t) : t \in [0,1]\}$).

If T is a trapezoid, let $T_0 = T \cap X_0$ and $T_1 = T \cap X_1$. In², Kaczynski proved the following result:

¹ Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

² Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

Theorem 1 *If T is a trapezoid, then .*

This theorem was used to construct a collection of non-intersecting paths (not lines) γ_x corresponding to each point $x \in [0,1]$ such that for each path $\lim_{t \rightarrow 0} \gamma_x(t) = (x,0)$ but the union satisfies

This construction immediately leads to the creation of a measurable half plane function with non-measurable boundary function³. We depict the situation in Figure 1.

One may wonder the extent to which this result holds if we omit the restriction that the lines of a trapezoid are not disjoint. To this end, we amend the trapezoid definition:

Definition 2 *An unrestricted trapezoid $T = \bigcup \gamma_x$ is a union (not necessarily disjoint) of line segments with each segment having its two endpoints on X_0 and X_1 (i.e. $\gamma_x = \{(1-t)x_0 + tx_1, t\} : t \in [0,1]$).*

In⁴, Kaczynski proves that there is no sensible upper bound in terms of $\mu_1(T_0)$ and $\mu_1(T_1)$ on the two dimensional measure of an unrestricted trapezoid.

Theorem 2 *There exists an unrestricted trapezoid T with $\mu_1(T_0) = \mu_1(T_1) = 0$ and $\mu_2(T) = \infty$.*

Proof: Let M be a residual set of measure zero (e.x. let x_k be an enumeration of rationals and $M = \bigcup \{x_k\}$)

). For any point (x,y) ($y \in (0,1)$) and line γ_x passing through (x,y)

with angle ϑ with the x -axis, let $F_0(\vartheta) = (x - y \cot(\vartheta), 0)$ and $F_1(\vartheta) = (x + (1 - y)\cot(\vartheta), 1)$ denote the intersections $\gamma_x \cap X_0$ and $\gamma_x \cap X_1$ respectively. F_0 and F_1 are homeomorphisms from $(0,\pi)$ onto X_0 and X_1 respectively, so $F_0^{-1}(M)$ and $F_1^{-1}(M)$ are both residual sets of measure zero. Since the intersection of residual

Figure 1: Plot of γ_x for $x \in [0,1]$ where $\gamma_{x_1} \cap \gamma_{x_2} = \emptyset$, $\lim_{t \rightarrow 0} \gamma_x(t) = (x,0)$, but $\mu_2(T) = \infty$. Given a nonmeasurable function $\varphi(x)$, if we define $f : (\mathbb{R} \times (0,\infty)) \rightarrow \mathbb{R}$ as $f(\gamma_x) = \varphi(x)$ and vanishing elsewhere, then f is measurable in the upper half plane but has a nonmeasurable boundary function φ .

sets is nonempty, for every (x,y) with $y \in (0,1)$ we can choose an angle) such that the line γ_x intersecting (x,y) with angle a intersects $X_0 \setminus M$ and $X_1 \setminus M$. Let the unrestricted trapezoid T be the union of all such lines so that $\mu_1(T_0) = \mu_1(T_1) = 0$ and $\mu_2(T) = \mu_2(\{(x,y), y \in (0,1)\}) = \infty$.

2. No lower bound

In⁵, Kaczynski poses the opposite question as an open problem, asking if there is a lower bound on the twodimensional measure of an unrestricted trapezoid in terms

³ Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

⁴ Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

⁵ Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

of $\mu(T_0)$ and $\mu(T_1)$. We will prove the following theorem indicating no lower bound exists:

Theorem 3 *There exists an unrestricted trapezoid T with $\mu_1(T_0) = \mu_1(T_1) = \square$ and $\mu_2(T) = 0$.*

Since it suffices to find an unrestricted trapezoid with measure zero and sets $\mu(T_0) = \mu(T_1) = \mu([0,1]) = 1$, we can formulate the question a different way:

Proposition 1 *There exists a bijection $f : [0,1] \rightarrow [0,1]$ such that, for the set $S = \int_0^1 ((1-t)x + tf(x), t) dt$, we have $\mu(S) = 0$.*

It is clear that Theorem 3 is a corollary of Proposition 1.

It is perhaps not immediate that such a function should exist. Any monotone f induces a standard trapezoid which abides by Theorem 1. The choice $f(x) = 1 - x$ gives measure $\mu(S) = 1/2$, and other decreasing functions cannot be massaged to induce a smaller measure. We will instead draw inspiration

Figure 2: Examples of the fractal construction of an unrestricted trapezoid with null measure. (Left) The base pattern induced by f_1 (Right) Illustration of unrestricted trapezoid induced by f .

from fractal geometry⁶; the particular iterative scheme we choose will create one-dimensional slices in x similar to the Cantor middle third set. In Figure 2, the basic pattern is represented by the piecewise function

otherwise we divide $[0,1]$ into three subintervals and connect them via parallelograms where f_1 essentially swaps the second and the third. In this case, $\mu(S) = 5/6$. In each subsequent iteration we embed an affine transformed version of the base pattern in each parallelogram, thus reducing the total measure of the unrestricted trapezoid (we will prove this limits to zero as in Proposition 1).

Since this procedure iteratively divides subintervals of $[0,1]$ in 3, it is perhaps not surprising that this this sequence of functions $\{f_k\}$ has a natural representation on base 3 fractions, namely

$$f_k(0.x_1x_2x_3\dots) = 0.y_1y_2\dots y_kx_{k+1}x_{k+2}\dots$$

where $y_j = 2x_j \pmod{3}$. Let $f = \lim_{k \rightarrow \infty} f_k$ such that for input $x \in [0,1]$, f replaces every 1 in its base 3 expansion with a 2 and vice-versa. We will prove that this function satisfies proposition 1. Before we prove proposition 1, we will state a result from⁷ on modified Cantor sets: **Proposition 2** *Let C_t be a modified Cantor set with representation*

Then the following holds:

⁶ Kenneth J. Falconer. *Techniques in fractal geometry*, volume 3. Chichester: Wiley, 1997.

⁷ Richard Kenyon. Projecting the one-dimensional sierpinski gasket. *Israel Journal of Mathematics*, 97:221–238, 1997.

in lowest terms, $p + q \not\equiv 0 \pmod{3}$

We are now ready to prove Proposition 1.

Proof of Proposition 1: Let us define f acting on base 3 fractions as

$$fk(0.x_1x_2x_3\dots) = 0.y_1y_2y_3\dots$$

where $y_j = 2x_j \pmod{3}$. Then the linear combination $(1 - t)x + tf(x)$ satisfies

Since St is just a scaled copy of $C(2-t)/(1+t)$, by Proposition 2 we have the measure in lowest terms, $p + q \not\equiv 0 \pmod{3}$

Since $\mu_1(St)$ takes on nonzero value at only countably many t , the result follows from Fubini's theorem. \square

3. Parallelogram formulation

Over 30 years after⁸, Kaczynski would outline a reformulation of the problem, offering three conjectures on the matter⁹. Rather than conceiving the unrestricted trapezoid as a union of lines, Kaczynski considered unions of increasingly thinner parallelograms.

Definition 3 *Let be a parallelogram with parallel sides $[(j - 1)/n, j/n] \times \{0\}$ and $[(k - 1)/n, k/n] \times \{1\}$. For $\sigma \in \text{Sym}(n)$, we say that the unrestricted trapezoid is given by the union.*

In this way, the unrestricted trapezoid associated with f_1 above can be expressed as.

The following conjectures center around properties of the smallest unrestricted trapezoid of n parallelograms of width $1/n$, namely

Conjecture 1 $\lim_{n \rightarrow \infty} a(n) = 0$.

Kaczynski claimed to have proved this. We will show that this follows from Proposition 1 in the following section.

Conjecture 2 *There exists constants $c_{\pm} > 0$ such that*

Kaczynski claimed to be able to prove that c_{-} exists and also claimed a weaker upper bound in the form of $a(n) < c_{+} (\log(\log n))^2 / \log n$. We will prove a different upper bound.

Conjecture 3 *The sequence $a(n)$ decreases monotonically.*

This was left as an open problem in¹⁰.

⁸ Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

⁹ Yahoo! News. The unabomber letters: A yahoo news special report, 2016. <https://www.yahoo.com/entertainment/the-unabomber-letters---a-yahoo-news-special-report-170846210.html>

¹⁰ Yahoo! News. The unabomber letters: A yahoo news special report, 2016. <https://www.yahoo.com/entertainment/the-unabomber-letters---a-yahoo-news-special-report-170846210.html>

4. Proving the conjectures

To prove conjectures from the previous section, we first connect projections of the Sierpinski gasket¹¹ to our modified Cantor sets.

Definition 4 We define the Sierpinski gasket $G \subset \mathbb{R}^2$ as

We define the n th partial Sierpinski gasket $G_n \subset G$ similarly. Let and let $\Delta(a,b,c)$ be the area bounded by the triangle with vertices $a,b,c \in \mathbb{R}^2$.

Let the projection operator $\text{proj}_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}$ map points $(x,y) \in \mathbb{R}^2$ onto a line through the origin with angle θ to the x -axis such that $\text{proj}_\theta : (x,y) \mapsto x\cos\theta + y\sin\theta$. We are interested in the sets $\text{proj}_\theta(G_n)$; more generally there is an active field of research surrounding linear projections of fractals in \mathbb{R}^2 ¹². One of the major subjects of interest is the following:

Definition 5 The Favard distance of a compact set $E \subset \mathbb{R}^2$ is given by

Fav

While it is perhaps obvious that $Fav(G) = 0$ from proposition 2, a result by Bond and Volberg¹³ gives a decay bound on the Favard distances of partial Sierpinski gaskets.

Proposition 3 There exist constants $C,p > 0$ such that $Fav(G_n) \leq Cn^{-p}$.

A later result by Bond and Volberg proves that $p > 1/14$ ¹⁴.

The rest of this section centers around utilizing proposition 3 to prove the conjectures from section 3. To do this, we must compare $\text{proj}_\theta(G_n)$ to the modified Cantor sets from Proposition 2 and the horizontal slices of the unrestricted trapezoid. Let us define partial variants of these sets:

Definition 6 Let T be the unrestricted trapezoid associated with the n th iteration induced by swapping 1 and 2 in a base 3 expansion (i.e. if $x = 1.x_1x_2\dots x_n$ is an n -digit base 3 number, then $\sigma(x) = 1.y_1y_2\dots y_n$ where $y_k = 2x_k \pmod{3}$), then we write $S(n,t) = (T \times \{t\}) \cap T\sigma^n$, the slice of, as

We similarly define the n th partial modified Cantor set as

This allows us to make the following comparison:

Lemma 1 For $t \in [0,1]$ and, we have

¹¹ Richard Kenyon. Projecting the one-dimensional sierpinski gasket. *Israel Journal of Mathematics*, 97:221–238, 1997.

¹² Izabella Laba. Recent progress on favard length estimates for planar cantor sets. In *Operator-Related Function Theory and Time-Frequency Analysis*, pages 117–145. Cham: Springer International Publishing.

¹³ Matthew Bond and Alexander Volberg. The power law for buffon’s needle landing near the sierpinski gasket. arXiv preprint arXiv:0911.0233, 2009.

¹⁴ Matthew Bond and Alexander Volberg. Buffon needle lands in ϵ -neighborhood of a 1-dimensional sierpinski gasket with probability at most $|\log\epsilon|^{-c}$. *Comptes Rendus. Mathématique*, 348(11-12):653–656, 2010.

Proof: We can show that for $t \in [0, 1]$ since

The result then follows from noticing that for, we have $\text{proj} \in [0, 1]$.

Combining these results leads us to a similar polynomial bound on the decay of the unrestricted trapezoids induced by the iterative scheme described in Definition 6.

Proposition 4 *Let σ be the permutation on base 3 numbers as defined in Definition 6. Then there exist constants $C, p > 0$ such that*

Proof: By Fubini's theorem, The measure of is given by integrating over the measures of the horizontal slices

From Lemma 1, this integral can be bound by an integral over projections of the Sierpinski gasket:

By the change of variables $\vartheta = \varphi(t)$, we can write:

$(\text{proj} \vartheta(G_n)) d\vartheta$ Placing an estimate on the trigonometric portion of the integral we can write

$(\text{proj}$

Since Lebesgue measure is nonnegative, we can extend the limits of integration to bound the measure of by a factor of the associated Favard distance

$\text{Fav}(G_n)$

Proposition 3 completes the proof. \square

This result translates to a logarithmic bound on $a(n)$ only for $n = 3m$ where $m \in \mathbb{N}$. If n has the base 3 representation $n = x_k x_{k-1} \dots x_0$, we can define σn as in Figure 3 such that the first $x_k 3^k$ intervals map to x_k copies of, the next $x_{k-1} 3^{k-1}$ intervals map to x_{k-1} copies of, etc. Using this scheme, we can prove a weaker upper bound on Conjecture 2 (consequently proving Conjecture 1 as well):

Theorem 4 *There exist constants $C, p > 0$ such that $a(n) < C/(\log n)^p$.*

Before proving this theorem, we supply a technical lemma:

Lemma 2 *For $p \in (0, 1)$, there exists a constant C such that*

Proof: Let us make the change of variables $t = 1 - x/n$ such that this integral can be transformed into

For any constant $C > 1$, we can find t_0 dependent only on C, p such that $pt + 1 > (1 - t)^{-p}$ for $t \in (0, t_0)$. This allows us to split the integral

Both of these integrals have closed form expressions, the leading asymptotic term $1/n$ arising from the left integral, thus completing the proof. \square

Proof of Theorem 4: It is clear that. Let $n = x_k x_{k-1} \dots x_0$ be a base 3 integer representation. Then the following holds:

Figure 3: Examples of parallelogram-defined unrestricted trapezoids where σ is as outlined in the proof of Theorem 4 (*Left*) $n = 1000$ which has base-3 representation 1101001 (*Right*) $n = 2000$ which has base-3 representation 2202002.

Each digit must satisfy $x_j \geq 2$, and by Proposition 4 there exist constants $C, p > 0$ such that, so we may write

Since the function $f(x) = 3xx - p$ is increasing for $x > p/\log 3(x)$, we can bound this sum by the integral

By making the change of variables $x = u \log 3 e$, we can write

The result follows from applying Lemma 2 and collecting constants. **5 Application to Cluster Sets**

While Theorem 1 assisted in constructing a measurable half-plane function with nonmeasurable boundary function in¹⁵, the open questions from¹⁶ and¹⁷ answered by Propositions 1 and 4 were also initially conceived to address problems related to boundary functions and cluster sets. We state the relevant material here for completeness.

Definition 7 Given a function defined on the open unit disk $f : D \rightarrow C$ and a Jordan arc γ with an endpoint on the boundary of the disk, we say that $C(f, \gamma)$ is the cluster set of f on γ where

$$C(f, \gamma) = \{w \in C : \exists \{z_n\} \text{ s.t. } z_n \in \gamma, |\lim z_n| = 1, \lim f(z_n) = w\}$$

Definition 8 Let $\gamma_1, \gamma_2, \gamma_3 \subset D$ be Jordan arcs with a similar endpoint z on the unit circle. If, for a disk function f , the intersection satisfies $C(f, \gamma_1) \cap C(f, \gamma_2) \cap C(f, \gamma_3) = \emptyset$, then f has the three-arc property at z . If each $\gamma_1, \gamma_2, \gamma_3$ is a union of rectilinear segments, then we say f has the three-segment property at z .

In¹⁸, a version of conjecture 1 was suggested to assist in answering the following open question from¹⁹ (presented as conjecture):

Conjecture 4 There exists a continuous function in D having the three-segment property at each point of a set of positive measure or second category on $|z| = 1$.

Weaker versions of this conjecture have been proved. Jarník²⁰ gave an example of a function having the three-segment property at uncountably many points. Piranian was cited in²¹ as having proved the existence of a continuous disk function with the three-arc (not segment) property. Bagemihl later proved the existence of a normal meromorphic (not continuous) disk function with the three-segment property²².

¹⁵ Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

¹⁶ Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

¹⁷ Yahoo! News. The unabomber letters: A yahoo news special report, 2016. <https://www.yahoo.com/entertainment/the-unabomber-letters---a-yahoo-news-special-report-170846210.html>

¹⁸ Theodore John Kaczynski. *Boundary Functions*. PhD thesis, University of Michigan, 1967.

¹⁹ Frederick Bagemihl, George Piranian, and G. S. Young. Intersections of cluster sets. *Bulletin of the Polytechnic Institute of Ias,i*, 5(9):29–34, 1959.

²⁰ Vojtěch Jarník. Sur les fonctions de deux variables réelles. *Fundamenta Mathematicae*, 27:147–150, 1936.

²¹ Frederick Bagemihl, George Piranian, and G. S. Young. Intersections of cluster sets. *Bulletin of the Polytechnic Institute of Ias,i*, 5(9):29–34, 1959.

²² Frederick Bagemihl. The three-arc and three-separated-arc properties of meromorphic functions. *Nagoya Mathematical Journal*, 53:137–140, 1974.

References

The Ted K Archive

A critique of his ideas & actions



Parker Kuklinski

An uncountable union of line segments with null two-dimensional measure
October 2023

[<arxiv.org/abs/2311.16210>](https://arxiv.org/abs/2311.16210)

See the attached file below and furthest right for the original PDF.

www.thetedkarchive.com